CAMBRIDGE INTERNATIONAL EXAMINATIONS

www.PapaCambridge.com **GCE Ordinary Level**

MARK SCHEME for the October/November 2013 series

5070 CHEMISTRY

5070/32

Paper 3 (Practical Test), maximum raw mark 40

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes should be read in conjunction with the question paper and the Principal Examiner Report for Teachers.

Cambridge will not enter into discussions about these mark schemes.

Cambridge is publishing the mark schemes for the October/November 2013 series for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level components and some Ordinary Level components.

Page 2	Mark Scheme	Syllabus	.0	V
	GCE O LEVEL – October/November 2013	5070	100	

(a) Titration

Accuracy (8 marks)

For the two best titres give:

- 4 marks for a value within 0.2 cm³ of supervisor 2 marks for a value within 0.3 cm³ of supervisor 1 mark for a value within 0.4 cm³ of supervisor

Concordance (3 marks)

Give:

- 3 marks if all the ticked values are within 0.2 cm³
- 2 marks if all the ticked values are within 0.3 cm³
- 1 mark if all the ticked values are within 0.4 cm³

Average (1 mark)

Give 1 mark if the candidate calculates a correct average (error not greater than 0.05) of all his/her ticked values. [12]

3Cambridge.com

			4	En.
Pag	ge 3	Mark Scheme	Syllabus	· 0
		GCE O LEVEL – October/November 2013	5070	123
Ass	uming a 2	25.0 cm ³ pipette and a titre of 25.2 cm ³ ,		WW. Patra Cambridge
(b)	moles of	f sulfuric acid present in average volume of Q		Topo
	= 25.2×0			
	1000)		
	= 0.0025	52		[1]
(c)	moles of	f sodium carbonate in P		
	$=\frac{25.0\times0}{100}$	0.02		
	100	0		
	= 0.0005	5		[1]
(d)	moles of	f sulfuric acid reacting with sodium carbonate		
	= 0.0005	5		[1]
(e)	moles of	f sulfuric acid reacting with sodium hydroxide		
	= 0.0025	52 – 0.0005		
	= 0.0020	02		[1]
(f)	concentr	ration of sodium hydroxide in P		
	=	$\frac{02 \times 2 \times 1000}{25.0}$		

[1]

[Total: 17]

 $= 0.162 \text{ mol/dm}^3$

			2.
Page 4	Mark Scheme	Syllabus	.0
	GCE O LEVEL – October/November 2013	5070	12

2 R is hydrochloric acid; S is sodium sulfite.

Observations		Notes		
General points For ppt/precipitate allow solid, suspension, powder				
For gases Name of gas requires test to be at least partially correct. effervesces = bubbles = gas vigorously evolved but not gas evolved				
For solutions colourless not equivalent to clear, clea	ar not equivale	ent to colourless		
Test 1				
white ppt	(1)			
Test 2				
insoluble in acid	(1)			
Test 3				
ppt disappears	(1)			
colourless solution	(1)			
Test 4				
effervescence	(1)			
turns limewater milky	(1)			
carbon dioxide	(1)	to score carbon dioxide mark there must be some indication of the limewater test e.g. 'tested with limewater'		
solid disappears	(1)			
Test 5				
effervescence	(1)			
pops with a lighted splint	(1)			
hydrogen	(1)	to score hydrogen mark there must be some indication of a test e.g. 'popped with a splint', 'tested with a burning splint'		
solid disappears	(1)			

Page 5	Mark Scheme	Syllabus	.0	V
	GCE O LEVEL – October/November 2013	5070	100	

Test 6			ambridg
turns colourless/decolourised (1)		(1)	23
Test 8			
(a)	white ppt	(1)	
(b)	ppt disappears	(1)	
	colourless solution	(1)	
Test 9			
(a)	white ppt	(1)	
(b)	ppt disappears	(1)	
(c)	coloured solid	(1)	if ppt remains in (b) allow mark in (c) providing the solid is not white
Test 10			
(a)	red solution	(1)	allow brown or orange or any mixture of these three colours
(b)	turns yellow	(1)	allow green or green-yellow or yellow-green
(c)	green or black ppt	(1)	do not allow any reference to brown e.g. black–brown
	insoluble in excess	(1)	

A cation present in **R** is hydrogen/H⁺ (bubbles or gas tested in test 4 or 5). [1]

An anion present in **R** is chloride/ Cl^- (tests 1 and 2 white ppt remains in acid). [1]

If cation and anion identifications are both correct but inverted allow 1 mark.

S is acting as a reducing agent/reductant. (in test 6 decolourised or green/black ppt in test 10)

[1]

Any 23 out of the 26 scoring points

[Total: 23]