



UNIVERSITY OF CAMBRIDGE INTERNATIONAL EXAMINATIONS
General Certificate of Education Ordinary Level

CANDIDATE NAME

CENTRE NUMBER

CANDIDATE NUMBER



CHEMISTRY **5070/22**
Paper 2 Theory **October/November 2013**
1 hour 30 minutes

Candidates answer on the Question Paper.
No Additional Materials are required.

READ THESE INSTRUCTIONS FIRST

Write your Centre number, candidate number and name on all the work you hand in.
Write in dark blue or black pen.
You may use a soft pencil for any diagrams, graphs or rough working.
Do not use staples, paper clips, highlighters, glue or correction fluid.
DO NOT WRITE IN ANY BARCODES.

Section A

Answer **all** questions.
Write your answers in the spaces provided in the Question Paper.

Section B

Answer any **three** questions.
Write your answers in the spaces provided in the Question Paper.

Electronic calculators may be used.
You may lose marks if you do not show your working or if you do not use appropriate units.

A copy of the Periodic Table is printed on page 16.
At the end of the examination, fasten all your work securely together.
The number of marks is given in brackets [] at the end of each question or part question.

This document consists of **16** printed pages.

Section A

Answer **all** the questions in this section in the spaces provided.

The total mark for this section is 45.

A1 Choose from the following elements to answer the questions below.

chlorine
hydrogen
iron
lithium
nickel
nitrogen
oxygen
potassium
silver
sulfur
vanadium
zinc

Each element can be used once, more than once or not at all.

Which element

- (a) is liberated at the anode when an aqueous solution of potassium sulfate is electrolysed,
..... [1]
- (b) is used as a catalyst in the manufacture of margarine,
..... [1]
- (c) is a non-metallic solid, an atom of which contains only six valency electrons,
..... [1]
- (d) is higher than sodium in the reactivity series,
..... [1]
- (e) is in Period 5 of the Periodic Table,
..... [1]
- (f) forms a white oxide which is amphoteric?
..... [1]

[Total: 6]

- A2** Carboxylic acids are a homologous series containing the $-\text{CO}_2\text{H}$ group.
The table shows some properties of the first four carboxylic acids in the series.

carboxylic acid	molecular formula	density in g/cm^3	boiling point in $^\circ\text{C}$
methanoic acid	CH_2O_2	1.220	101
	$\text{C}_2\text{H}_4\text{O}_2$	1.049	118
propanoic acid	$\text{C}_3\text{H}_6\text{O}_2$	0.993	141
butanoic acid	$\text{C}_4\text{H}_8\text{O}_2$	0.958	165

- (a) (i) Describe how the density of these carboxylic acids varies with the number of carbon atoms in the molecule.

..... [1]

- (ii) Name the carboxylic acid with the molecular formula $\text{C}_2\text{H}_4\text{O}_2$.

..... [1]

- (iii) Draw the structure of propanoic acid, showing all atoms and bonds.

[1]

- (b) The next carboxylic acid in this homologous series is pentanoic acid.
Pentanoic acid has five carbon atoms.

- (i) Deduce the molecular formula for pentanoic acid.

..... [1]

- (ii) Suggest a value for the boiling point of pentanoic acid.

..... $^\circ\text{C}$ [1]

- (c) Butanoic acid, $\text{C}_3\text{H}_7\text{CO}_2\text{H}$, reacts with sodium to form a salt and a gas.

- (i) Name the gas.

..... [1]

- (ii) Give the formula of the salt.

..... [1]

- (d) Esters are formed when carboxylic acids react with alcohols.
The reaction is catalysed by hydrogen ions.


(i) Describe and explain the effect of a catalyst on reaction rate.

.....
.....
..... [2]

(ii) State one commercial use of esters.

..... [1]

(iii) The structure of an ester is shown below.



Name this ester.

..... [1]

[Total: 11]

A3 Silicon is an element in Group IV of the Periodic Table.

(a) Give the electronic configuration for a silicon atom.

..... [1]

(b) Silicon has three naturally occurring isotopes.

Complete the following table for two of these isotopes.

isotope	^{28}Si	^{30}Si
number of protons		
number of electrons		
number of neutrons		

[3]

(c) Silicon reacts with chlorine on heating to form silicon(IV) chloride, SiCl_4 .

Construct an equation for this reaction.

[1]

(d) Silicon(IV) chloride is a simple molecular compound.

(i) Suggest **two** physical properties of silicon(IV) chloride other than solubility.

1


2 [2]

(ii) Draw a 'dot-and-cross' diagram for silicon(IV) chloride.

You only need to show the outer shell electrons for each atom.

[2]

- (e) Silicon(IV) chloride reacts with water to form silicon(IV) oxide. Part of the structure of silicon(IV) oxide is shown below.

**Key**

- silicon atom
○ oxygen atom

Explain, in terms of structure and bonding, why silicon(IV) oxide has a very high melting point.

.....

.....

.....

..... [2]

[Total: 11]

A4 The carbon cycle regulates the amount of carbon dioxide in the atmosphere.

(a) (i) State **two** processes which release carbon dioxide into the atmosphere.

1

2 [2]

(ii) Name one process which removes carbon dioxide from the atmosphere.

..... [1]

(b) Carbon dioxide is a greenhouse gas.

(i) What is the meaning of the term *greenhouse gas*?

.....


..... [1]

(ii) Name another greenhouse gas and give a natural source of this gas.

name

source [2]

(c) Carbon dioxide dissolves in water to form a weakly acidic solution.



(i) What is the meaning of the term *weak acid*?

.....

..... [1]

(ii) Describe how you could measure the pH of this solution other than by using a pH meter.

.....

.....

..... [2]

(d) Sodium hydrogencarbonate, NaHCO_3 , decomposes on heating to form a carbonate, water and a gas which turns limewater milky. Construct an equation for this reaction.

[2]

[Total: 11]

A5 A student reacts magnesium ribbon with excess hydrochloric acid. She follows the course of the reaction by measuring the volume of gas produced against time.

(a) Write the equation for the reaction of magnesium with hydrochloric acid.

.....[1]

(b) (i) On the axes below draw a sketch graph to show how the volume of gas produced during the reaction varies with time and label this line 'A'. Label the axes with the appropriate units.



[2]

(ii) The student then carries out the experiment at a **lower** temperature. All the other conditions remain the same.

On the axes above draw another line to show how the volume of gas produced varies with time and label this line 'B'. [1]

(c) Magnesium reacts with carbon to form the compound magnesium carbide.

Calculate the percentage by mass of magnesium in magnesium carbide, MgC_2 .

[2]

[Total: 6]

Section B

Answer **three** questions from this section in the spaces provided.

The total mark for this section is 30.

B6 Aluminium is extracted from purified bauxite by electrolysis.

- (a) Describe how this electrolysis is carried out and construct equations for the reactions occurring at both the anode and cathode.

.....

 [4]

- (b) What properties of aluminium make it useful for


- (i) making aircraft,
 (ii) making electricity cables. [2]

- (c) Aluminium is high in the reactivity series.

- (i) Explain why aluminium does not react with aqueous copper(II) sulfate.

.....
 [2]

- (ii) When a few drops of aqueous sodium chloride are added to a mixture of aluminium and aqueous copper(II) sulfate, a vigorous reaction occurs.



What type of reaction is this?

..... [1]

- (iii) State the formula of aluminium sulfate.

..... [1]

[Total: 10]

B7 Ethene is an unsaturated hydrocarbon.

(a) What is the meaning of each of these terms?

unsaturated

hydrocarbon [2]

(b) Ethene can be manufactured by cracking.

(i) State the conditions used for cracking.

.....
 [2]

(ii) Construct an equation for the cracking of tetradecane, $C_{14}H_{30}$, to form ethene and one other hydrocarbon.


..... [1]

(c) Alkenes such as ethene can undergo addition polymerisation.

(i) State one use of poly(ethene).

..... [1]


(ii) The diagram below shows a section of a polymer chain.



Deduce the structure of the monomer which is used to make this polymer.

[1]

- (d) Ethanol can be manufactured by the catalytic addition of steam to ethene.



If the reactants are not recycled, only 5% of the ethene is converted to ethanol.


Calculate the mass of ethanol formed from 0.4 tonnes of ethene when only 5% of the ethene is converted to ethanol.

[1 tonne is 1 000 000 grams]

mass of ethanol tonnes [3]

[Total: 10]

- B8** When hydrogen iodide, HI, is heated in a closed tube, the following dynamic equilibrium is established.



- (a) What is meant by the term *dynamic equilibrium*?

.....

 [2]

- (b) The table shows the concentrations of HI(g), H₂(g) and I₂(g) in the equilibrium mixture at 25 °C and 450 °C.

substance	concentration at 25 °C / mol/dm ³	concentration at 450 °C / mol/dm ³
HI(g)	0.94	0.79
H ₂ (g)	0.033	0.11
I ₂ (g)	0.033	0.11

- (i) The tube has a volume of 50 cm³.

Calculate the mass of hydrogen iodide in the equilibrium mixture at 25 °C.


[2]

- (ii) Describe and explain the differences in the concentrations of reactant and products at 25 °C and 450 °C.

.....

 [2]

- (c) Complete the energy profile diagram for the decomposition of hydrogen iodide. On your diagram label
- the products,
 - the enthalpy change of the reaction, ΔH .



[2]

- (d) An aqueous solution of hydrogen iodide contains iodide ions.

Describe a test for iodide ions.

.....
.....

[2]

[Total: 10]

B9 The compounds ammonium nitrate and ammonium sulfate are both fertilisers.

(a) Explain why farmers add these fertilisers to soils.

.....
 [1]

(b) Ammonium sulfate can be prepared by adding sulfuric acid to aqueous ammonia.

Construct the equation for this reaction.

..... [1]

(c) Excess acidity in soils can be treated by adding calcium hydroxide.

(i) Give the formula of the ion present in calcium hydroxide which causes it to be alkaline.


..... [1]

(ii) Explain why adding calcium hydroxide causes loss of nitrogen from fertilisers such as ammonium nitrate, which have been previously added to the soil.

.....

 [2]


(d) A student titrated 10.0 cm³ of aqueous calcium hydroxide with hydrochloric acid.



It required 4.00 cm³ of 0.0100 mol/dm³ hydrochloric acid to neutralise 10.0 cm³ of aqueous calcium hydroxide.

Calculate the concentration of the calcium hydroxide.

..... mol/ dm³ [3]



(e) Describe how to obtain pure dry crystals of calcium chloride from an aqueous solution of calcium chloride.

.....

.....

.....

.....

..... [2]

[Total: 10]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

University of Cambridge International Examinations is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.

DATA SHEET
The Periodic Table of the Elements

		Group																																																																																															
I	II	III	IV	V	VI	VII	0																																																																																										
7 Li Lithium 3	9 Be Beryllium 4	1 H Hydrogen 1	11 B Boron 5	12 C Carbon 6	13 Al Aluminium 13	14 Si Silicon 14	15 P Phosphorus 15	16 S Sulfur 16	17 Cl Chlorine 17	18 Ar Argon 18	19 F Fluorine 9	20 Ne Neon 10	2 He Helium 2																																																																																				
23 Na Sodium 11	24 Mg Magnesium 12	27 Co Cobalt 27	28 Ni Nickel 28	29 Cu Copper 29	30 Zn Zinc 30	31 Ga Gallium 31	32 Ge Germanium 32	33 As Arsenic 33	34 Se Selenium 34	35 Br Bromine 35	36 Kr Krypton 36	37 Rb Rubidium 37	38 Sr Strontium 38	39 Y Yttrium 39	40 Ca Calcium 20	41 Nb Niobium 41	42 Mo Molybdenum 42	43 Tc Technetium 43	44 Ru Ruthenium 44	45 Rh Rhodium 45	46 Pd Palladium 46	47 Ag Silver 47	48 Cd Cadmium 48	49 In Indium 49	50 Sn Tin 50	51 Sb Antimony 51	52 Te Tellurium 52	53 I Iodine 53	54 Xe Xenon 54	55 Cs Caesium 55	56 Ba Barium 56	57 La Lanthanum 57	58 Ce Cerium 58	59 Pr Praseodymium 59	60 Nd Neodymium 60	61 Pm Promethium 61	62 Sm Samarium 62	63 Eu Europium 63	64 Gd Gadolinium 64	65 Tb Terbium 65	66 Dy Dysprosium 66	67 Ho Holmium 67	68 Er Erbium 68	69 Tm Thulium 69	70 Yb Ytterbium 70	71 Lu Lutetium 71	72 Fr Francium 87	88 Ra Radium 88	89 Ac Actinium 89	90 Th Thorium 90	91 Pa Protactinium 91	92 U Uranium 92	93 Np Neptunium 93	94 Pu Plutonium 94	95 Am Americium 95	96 Cm Curium 96	97 Bk Berkelium 97	98 Cf Californium 98	99 Es Einsteinium 99	100 Fm Fermium 100	101 Md Mendelevium 101	102 No Nobelium 102	103 Lr Lawrencium 103	133 Cs Caesium 55	137 Ba Barium 56	139 La Lanthanum 57	140 Ce Cerium 58	141 Pr Praseodymium 59	144 Nd Neodymium 60	147 Pm Promethium 61	150 Sm Samarium 62	152 Eu Europium 63	157 Gd Gadolinium 64	159 Tb Terbium 65	162 Dy Dysprosium 66	165 Ho Holmium 67	167 Er Erbium 68	169 Tm Thulium 69	173 Yb Ytterbium 70	175 Lu Lutetium 71	223 Fr Francium 87	226 Ra Radium 88	227 Ac Actinium 89	232 Th Thorium 90	231 Pa Protactinium 91	238 U Uranium 92	237 Np Neptunium 93	244 Pu Plutonium 94	243 Am Americium 95	247 Cm Curium 96	247 Bk Berkelium 97	251 Cf Californium 98	252 Es Einsteinium 99	257 Fm Fermium 100	258 Md Mendelevium 101	259 No Nobelium 102	260 Lr Lawrencium 103

* 58–71 Lanthanoid series
† 90–103 Actinoid series

Key

a	X	a = relative atomic mass
	X	X = atomic symbol
b		b = atomic (proton) number

The volume of one mole of any gas is 24 dm³ at room temperature and pressure (r.t.p.).